

GROUPS WITH MAXIMAL IRREDUNDANT COVERS AND MINIMAL BLOCKING SETS

ALIREZA ABDOLLAHI

ABSTRACT. Let n be a positive integer. Denote by $\text{PG}(n, q)$ the n -dimensional projective space over the finite field $\mathbb{F}_q$ of order q . A blocking set in $\text{PG}(n, q)$ is a set of points that has non-empty intersection with every hyperplane of $\text{PG}(n, q)$. A blocking set is called minimal if none of its proper subsets are blocking sets. In this note we prove that if $\text{PG}(n_i, q)$ contains a minimal blocking set of size k_i for $i \in \{1, 2\}$, then $\text{PG}(n_1 + n_2 + 1, q)$ contains a minimal blocking set of size $k_1 + k_2 - 1$. This result is proved by a result on groups with maximal irredundant covers.

1. Introduction and Results

Let G be a group. A set $\mathcal{S}$ of proper subgroups of G is called a cover for G whenever $G = \bigcup_{H \in \mathcal{S}} H$. The cover $\mathcal{S}$ is called irredundant if no proper sub-collection of $\mathcal{S}$ is a cover for G . The cover $\mathcal{S}$ is called an n -cover if $|\mathcal{S}| = n$. The cover $\mathcal{S}$ is called maximal if each member of $\mathcal{S}$ is a maximal subgroup of the group G . The cover $\mathcal{S}$ is called core-free if the core of $D = \bigcap_{H \in \mathcal{S}} H$ in G is trivial, i.e. $D_G = \bigcap_{g \in G} g^{-1} D g$ is the trivial subgroup of G . The cover $\mathcal{S}$ for a group is called a $\mathfrak{C}_n$ -cover if $\mathcal{S}$ is a maximal irredundant core-free n -cover.

Let n be a positive integer. Denote by $\text{PG}(n, q)$ the n -dimensional projective space over the finite field $\mathbb{F}_q$ of order q . A blocking set in $\text{PG}(n, q)$ is a set of points that has non-empty intersection with every hyperplane of $\text{PG}(n, q)$. A blocking set is called minimal if none of its proper subsets are blocking sets.

There is a well-known relationship between minimal blocking sets in $\text{PG}(n, q)$ and irredundant covers of the abelian group $\mathbb{F}_q^{n+1}$, the direct product of $n+1$ copies of $\mathbb{F}_q$ (see e.g., [1]). In fact a minimal blocking set of size k exists in $\text{PG}(n, q)$ if and only if an irredundant k -cover for $\mathbb{F}_q^{n+1}$ whose all members are hyperplane exists (see e.g., Proposition 2.2 of [1]).

The main results of this note are the following. Theorem 1.1 is inspired by the proof of Lemma 3.1 of [3].

Theorem 1.1. *Let $\mathcal{C}_1 = \{M_1, \dots, M_m\}$ and $\mathcal{C}_2 = \{N_1, \dots, N_n\}$ be irredundant m - and n -covers for two groups G_1 and G_2 , respectively. Suppose that M_1 and N_1 are normal subgroups of G_1 and G_2 , respectively; and assume that a and b are elements of $G_1 \setminus M_1$ and $G_2 \setminus N_1$, respectively, such that $M_1 a$ and $N_1 b$ have the same order p for some prime number p . Then*

$$\mathcal{C} = \{(M_1 \times N_1)\langle(a, b)\rangle, M_i \times G_2, G_1 \times N_j \mid i = 2, \dots, m; j = 2, \dots, n\}$$

1991 *Mathematics Subject Classification.* 20D60; 51E21.

Key words and phrases. Maximal irredundant covers; minimal blocking sets .

is an irredundant $(n + m - 1)$ -cover for $G_1 \times G_2$ with intersection $D_1 \times D_2$, where $D_1 = \cap_{i=1}^m M_i$, $D_2 = \cap_{j=1}^n N_j$ and $(M_1 \times N_1)\langle(a, b)\rangle$ is the subgroup of $G_1 \times G_2$ generated by $M_1 \times N_1$ and the element (a, b) . In particular, if both $\mathcal{C}_1$ and $\mathcal{C}_2$ are maximal, then $\mathcal{C}$ is a maximal cover for $G_1 \times G_2$.

Corollary 1.2. *Let $\text{PG}(n_i, q)$ contain a minimal blocking set of size k_i for $i \in \{1, 2\}$. Then $\text{PG}(n_1 + n_2 + 1, q)$ contains a minimal blocking set of size $k_1 + k_2 - 1$.*

2. Proofs

Proof of Theorem 1.1. It is clear that $\mathcal{C}$ is an $(m + n - 1)$ -cover for $G_1 \times G_2$. We now prove that $\mathcal{C}$ is irredundant. First, $(M_1 \times N_1)\langle(a, b)\rangle$ is an irredundant member of $\mathcal{C}$: for $\mathcal{C}_1$ and $\mathcal{C}_2$ are both irredundant, there exist elements $x \in M_1 \setminus \bigcup_{i=2}^m M_i$ and $y \in N_1 \setminus \bigcup_{j=2}^n N_j$. It follows that the element (x, y) , among the members of $\mathcal{C}$, belongs only to $(M_1 \times N_1)\langle(a, b)\rangle$ and so $(M_1 \times N_1)\langle(a, b)\rangle$ is an irredundant member. Now we show that each $M_i \times G_2$ is an irredundant member for the cover (for each $i = 2, \dots, m$). Without loss of generality and for convenience, we prove only the latter statement for $i = 2$, the other cases are similar. Let $x_0 \in M_2 \setminus \bigcup_{i \neq 2}^m M_i$ and $y_0 \in N_1 \setminus \bigcup_{j=2}^n N_j$. Then (x_0, y_0) , among the members of $\mathcal{C}$, could only possibly belong to $(M_1 \times N_1)\langle(a, b)\rangle$. If it is possible, then $x_0 \in M_1 a^\ell$ and $y_0 \in N_1 b^\ell$ for some integer ℓ . Since $x_0 \notin M_1$, $a^\ell \notin M_1$ and so p does not divide ℓ . This implies that $b^\ell \notin N_1$. Thus $y_0 \in N_1 \cap N_1 b^\ell = \emptyset$, a contradiction. Thus $M_2 \times G_1$ is an irredundant member of $\mathcal{C}$. By a similar argument, one may prove that each $G_1 \times N_j$ is an irredundant member for the cover (for each $j = 2, \dots, n$). This completes the proof of irredundancy of the cover $\mathcal{C}$.

Now by Lemma 2.2(b) of [2], $D_1 = \bigcap_{i=2}^m M_i$, $D_2 = \bigcap_{j=2}^n N_j$ and

$$\bigcap_{S \in \mathcal{C}} S = \left(\bigcap_{i=2}^m M_i \times G_2 \right) \bigcap \left(\bigcap_{j=2}^n G_1 \times N_j \right).$$

It follows that the intersection of the cover $\mathcal{C}$ is $D_1 \times D_2$.

For the last statement, note that it follows from the hypothesis that $|G_1 : M_1| = |G_2 : N_1| = p$, as we are assuming both M_1 and N_1 are maximal. Thus $(M_1 \times N_1)\langle(a, b)\rangle$ has prime index p in $G_1 \times G_2$ and so it is a maximal subgroup of $G_1 \times G_2$. The other members of the cover $\mathcal{C}$ are clearly maximal subgroups of $G_1 \times G_2$. This completes the proof of the last statement. $\square$

Corollary 2.1. *Let $V(n_i, q)$ ($i = 1, 2$) be the n_i -dimensional vector space over the finite field of order q . If $\mathcal{C}_1 = \{M_1, \dots, M_m\}$ and $\mathcal{C}_2 = \{N_1, \dots, N_n\}$ be irredundant m - and n -covers for the abelian groups $V(n_1, q)$ and $V(n_2, q)$, respectively; and assume that a and b are arbitrary elements of $G_1 \setminus M_1$ and $G_2 \setminus N_1$, respectively. Then*

$\mathcal{C} = \{(M_1 \times N_1)\langle(a, b)\rangle, M_i \times V(n_2, q), V(n_1, q) \times N_j \mid i = 2, \dots, m; j = 2, \dots, n\}$ is an irredundant $(n + m - 1)$ -cover for $V(n_1, q) \times V(n_2, q)$ with intersection $D_1 \times D_2$, where $D_1 = \cap_{i=1}^m M_i$ and $D_2 = \cap_{j=1}^n N_j$. In particular, if all members of $\mathcal{C}_1$ and $\mathcal{C}_2$ are hyperplanes of the corresponding spaces, then all members of $\mathcal{C}$ are hyperplanes of $V(n_1, q) \times V(n_2, q)$.

Proof of Corollary 1.2. It follows from Proposition 2.2 of [1] and Corollary 2.1. $\square$

Acknowledgement. The author is grateful to the referee for his/her helpful comments. This research was in part supported by a grant from IPM (No. 87200118). This research was also partially supported by the Center of Excellence for Mathematics, University of Isfahan.

REFERENCES

- [1] A. Abdollahi, M.J. Ataei and A. Mohammadi Hassanabadi, Minimal blocking sets in $PG(n, 2)$ and covering groups by subgroups, to appear in *Comm. Algebra*, **36** (2008).
- [2] R. A. Bryce, V. Fedri, L. Serena, Covering group with subgroups. *Bull. Austral. Math. Soc.*, **55** (1997) 469-476.
- [3] M. J. Tomkinson, Groups covered by finitely many cosets or subgroups, *Comm. Algebra*, **15** (1987) No. 4, 845-859.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ISFAHAN, ISFAHAN 81746-73441, IRAN; AND
SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), P.O.Box:
19395-5746, TEHRAN, IRAN. E-MAIL: abdollahi@member.ams.org a.abdollahi@math.ui.ac.ir